

BATS IN THE DIET OF OWLS FROM THE RUSSIAN FAR EAST, SOUTHERN SIKHOTE ALIN

VALENTINA V. ROSINA^{1*}, VALERIY P. SHOKHRIN²

¹Borissiak Paleontological Institute RAS, Profsovnaya str. 123, Moscow, Russia, 117997

*Corresponding author, email: ros@paleo.ru

²Lazovsky State Nature Reserve, Centralnaya str. 56, Lazo, Primorskii Region, Russia, 692890

Received 15 September 2010; accepted 30 May 2011

ABSTRACT - The pellets and nesting litter of six owl species from Southern Sikhote Alin were examined for bat remains. *Bubo bubo*, *Asio otus*, *A. flammeus* and *Strix uralensis* were found to prey occasionally on bats. Remains of four bat species - *Vespertilio murinus*, *Murina hilgendorfi*, *M. ussuriensis* and *Myotis* cf. *petax* were found. Owls preyed on bats more frequently in autumn and spring, during seasonal migrations of bats and when young animals are probably more abundant.

Key words: Chiroptera, Strigiformes, predation, feeding habits, Russia

RIASSUNTO - *I chiroterteri nella dieta di Strigiformi dell'estrema Russia orientale, Sikhote Alin meridionale*. Le borre e i resti alimentari di nidiate di sei Strigiformi, raccolti nel Sikhote Alin meridionale, erano esaminati per la ricerca di pipistrelli. *Bubo bubo*, *Asio otus*, *A. flammeus* and *Strix uralensis* predavano occasionalmente le seguenti 4 specie: *Vespertilio murinus*, *Murina hilgendorfi*, *M. ussuriensis* e *Myotis* cf. *petax*. La predazione sui pipistrelli era più frequente durante la stagione migratoria (autunno e primavera), quando i giovani sono probabilmente più abbondanti.

Parole chiave: Chiroterteri, Strigiformi, predazione, dieta, Russia

DOI: 10.4404/Hystrix-22.1-4503

INTRODUCTION

Bat populations of South Sikhote Alin are abundant and diverse (Rossina 2005; 2007), representing a major component of ecosystems. Being active at night, bats might be preyed by nocturnal predators (Kuzyakin 1950; Ruprecht 1990; 2005; Nowak 1994; Kowalski 1995). Owls rarely prey upon bats in the Far East of Russia. However, bats have been reported to be preyed on there by hawk owls *Ninox*

scutulata (Pukinskii 2003) and Ural owls *Strix uralensis* (Shibnev 1989). As Strigiformes are particularly abundant in Southern Sikhote Alin, mainly in the Lazovsky State Nature Reserve (LSNR; Vorob'ev 1954; Belopol'skii 1955; Shokhrin 2009), we investigated the role of six owl species in the Primorskii Region - eagle owl *Bubo bubo*, fish owl *Ketupa blakistoni*, long-eared owl *Asio otus*, marsh owl *A. flammeus*, oriental scops-owl *Otus sunia* and Ural owl *S. uralensis* - as bat predators. In

addition, available data on owls preying bats are discussed.

MATERIAL AND METHODS

The material examined was collected in the LSNR and some neighbouring areas (particularly in surroundings of Lazo village and near the town of Partizansk). Some pellets were found in the Mis Hill area in the valley of the River Kievka, about 10 km east of Lazo village and on Opasniy Island (in the northern part of the Sea of Japan, next to the northern border of the LSNR).



Figure 1 - Study area: Southern Sikhote Alin in the Russian Far East and Lazovsky State Nature Reserve (1).

The LSNR is located in the south-eastern part of the Primorskii Region at the foot of the Sikhote Alin range of mountains, facing the Sea of Japan (Fig. 1), and bordered by the valley of the rivers Kievka and Chernaya. The south-eastern end of the reserve stretches along the sea coast, including the small islands of Petrov and Belzov. At present the LSNR covers about 121000 ha and

is surrounded by a 15000 ha wide additional protected area. The total length of the border is 240 km, of which 36 km are along the seacoast. The average altitude is 500-700 m a.s.l., with some mountain peaks reaching 1400 m (Khokhryakov and Shokhrin 2002).

Food remains of *B. bubo* were collected in April-May and October-November 2003-2006, and in June 2005 and 2007-2008 on Petrov Island. Some pellets were collected on Opasniy Island and near Partizansk in April-May 2000-2001. Food remains of *A. otus* were collected in the LSNR in May-June 2001, 2004, 2005 and 2008; those of *A. flammeus* in October 2003 and May-June 2005; those of *K. blakistoni* in April 2001, while those of *S. uralensis* were collected in April-May 2000-2006 and 2008; finally, materials from *O. sunia* were collected in May 2005 and July 2006.

In practice, complete sets of bat skeleton bones can be found in owl pellets. The phalanges, forearms, mandibles, humeri and femurs are the most numerous and abundant remains. The pelvis and scapulae are also quite common, while tibias, maxillae and cranium fragments are found only occasionally. Intact craniums of bats are rather common.

The richness of bat bones in pellets and nesting litter was expressed by the following index: $R = (N \times 100) / \Sigma n$, where N = total number of bat remains and Σn = number of pellets and nesting litters with bats. In addition, the minimum number of bat individuals (Min) was calculated based on the number of the right and left bat skeletal fragments found in each pellet or nesting litter. Seasonal variation in both the number of pellets containing bat remains and the minimum number of preyed individuals was tested by the chi-squared test (χ^2), using STATISTICA, version 6.0 (StatSoft Inc. 2001).

The feeding habits of owls were assessed by the analysis of a total of 1466 pellets, 252 leftovers and 19 nesting litters (Tab. 1).

Table 1 - Bat remains in the owl materials collected in the Southern Sikhote Alin; P: pellets with bat remains; NI: nesting litters with bat remains; L: leftovers.

	Spring			Summer			Autumn
	P	NI	L	P	NI	L	P
Strigiformes							
<i>Bubo bubo</i>	79 (10)			85 (3)	2 (2)	104	53 (3)
<i>Asio otus</i>	512 (15)	11 (3)	69	40	1		
<i>Strix uralensis</i>	596 (2)	3					
<i>Asio flammeus</i>				55		14	7 (2)
<i>Ketupa blakistoni</i>	10						
<i>Otus sunia</i>	8			21	2	65	
Sum	1205 (27)	14 (3)	69	201 (3)	5 (2)	183	60 (5)
% bat remains	2.24	21.43	0	1.49	40	0	8.33
% bats in total food remains	2.33			1.28			8.33

RESULTS AND DISCUSSION

Bats were recorded in the food remains of four owl species: *B. bubo*, *A. otus*, *A. flammeus* and *S. uralensis*. The literature indicates that *K. blakistoni* mainly preys on fish and amphibians (Pukinskii 1977; 2003; Voronov and Zdorikov 1988), while insects and arachnids have been reported to be the main food of *O. sunia* (Pukinskii 2003; Shokhrin 2009). To our knowledge, bats have never been found in the diet of these owls. However, some bat remains were found in the pellets of pygmy owl *Glaucidium passerinum* and Tengmalm's owl *Aegolius funereus* (Shepel 1992), which are only slightly larger than *O. sunia*.

Four bat species, *i.e.* parti-coloured bat *Vespertilio murinus*, Eastern water bat *Myotis cf. petax*, Siberian tube-nosed bat *Murina hilgendorfi* and Ussuri tube-nosed bat *M. ussuriensis*, were found to be preyed upon.

Overall, bat remains were found in 35 pellets (2.38%) and 5 nesting litters (26.31%; Tab. 1), *i.e.* only 2.3% of the materials examined. Pellets with bat remains were more frequent in autumn ($\chi^2 = 7.59$, $P = 0.022$; Tab. 1), while the minimum number of bats recorded was the highest in spring ($\chi^2 = 72.3$, $P < 0.0001$; Tab. 2), corresponding to the periods of migration of some species of bats to and back from hibernation roosts. The highest concentration of bat bones was recorded in autumn pellets, ($R = 51.4\%$; Tab. 2), which mostly included bones of subadult bats, whilst the maximum bat species diversity in the food remains of owls was recorded in spring (Tab. 2). In summer pellets with bats were rarely found as, being mostly washed off by summer rains, they tended to disintegrate.

The eagle owl preyed on bats more frequently than other owls (Tab. 1). On the whole, 90% of pellets with bats belonged to *B. bubo* and *A. otus*. *B. bubo* is

Table 2 - Seasonal distribution of Chiroptera remains in pellets and nesting litter of Strigiformes in Southern Sikhote Alin; *n*: number of pellets and nesting litters with bat remains; *N*: total number of bat bones in pellets and nesting litters; *Min*: minimum number of bat individuals; *R*: index of richness of bat bones in pellets and nesting litters; *Bb*: *Bubo bubo*; *Ao*: *Asio otus*; *Su*: *Strix uralensis*; *Af*: *Asio flammeus*.

		Spring				Summer		Autumn		
Chiroptera		<i>Bb</i> <i>n</i> =10	<i>Ao</i> <i>n</i> =15+3	<i>Su</i> <i>n</i> =2	%	<i>Bb</i> <i>n</i> =3+2	%	<i>Bb</i> <i>n</i> =3	<i>Af</i> <i>n</i> =2	%
<i>V. murinus</i>	N	147	-	-	-	56	-	240	17	-
	Min	21	-	-	40.4	6	85.7	12	2	100
<i>M. hilgendorfi</i>	N	-	60	-	-	-	-	-	-	-
	Min	-	4	-	7.7	-	-	-	-	-
<i>M. ussuriensis</i>	N	-	284	85	-	-	-	-	-	-
	Min	-	18	5	44.2	-	-	-	-	-
<i>M. cf. petax</i>	N	1	-	-	-	1	-	-	-	-
	Min	1	-	-	1.9	1	14.3	-	-	-
Chiroptera indet.	N	2	1	-	-	-	-	-	-	-
	Min	2	1	-	5.8	-	-	-	-	-
R		19.33				11.4		51.4		
Location		Island Petrov	Lazo, Island Petrov	Kievka River Valley		Island Petrov, Kievka River Valley		Island Petrov		

a generalist predator (Shepel 1992). In the Primorskii Region the diet of *B. bubo* includes approximately 60 species of vertebrates: 31.1-98.0% of mammals; 2.0-68.9% of birds; 0.9-5.0% of amphibians (Tab. 3; see also Shokhrin 2005a).

Bat remains are found quite frequently in the pellets of *B. bubo* (Obuch 1989; Stubbe et al. 1989; Ruprecht 1990; 2005), and were also previously recorded in the Primorskii Region (Kolomiitsev and Poddubnaya 1985). On Petrov Island, we found two bat species to be preyed upon by this owl: *V. murinus* and *Myotis cf. petax* (Tab. 2). Remains of *V. murinus* were very abundant in autumn pellets (Tab. 2), which

interestingly were all collected on the sea shore. Probably, the eagle-owl hunts these bats during their seasonal spring and autumn migrations along the coast. During these seasons, some eagle-owl pellets consisted solely of bat bones and hair, representing up to 7 specimens of *V. murinus* (Tab. 2). About 50% of bat bones in autumn pellets belonged to subadults, which probably are less able to escape predation.

In autumn *V. murinus* was also preyed by *A. flammeus* (Tab. 2) for which murine rodents are the main food resources in the Primorskii Region (Tab. 3; Shepel 1992; Priklonskii 1993; Shokhrin 2005a, b; 2009).

Table 3 - Diet of four owl species from Southern Sikhote Alin in 2000-2008.

Prey groups (in %)	<i>Bubo bubo</i>	<i>Asio otus</i>	<i>Asio flammeus</i>	<i>Strix uralensis</i>
Mammals:	63.7	96.2	92.9	92.7
Muroidea	53.5	93.3	90.8	84.0
Other Rodentia	3.7	0.4	0.5	3.1
Mustelidae	0.4	0.1	-	0.1
Insectivora	0.1	0.8	-	5.2
Chiroptera	5.9	1.6	1.6	0.3
Aves	34.2	2.3	5.5	5.9
Reptilia	-	0.1	-	-
Amphibia	2.2	1.0	1.6	1.3
Insect	0.1	0.4	-	0.1
Total number of specimens	763	1717	184	1983

Bats are also included in the diet of marsh owls in the Ulyanovsk Region (Zhitkov and Buturlin 1906) and in some regions of Finland (0.2% of all owl preys; Aho 1964).

S. uralensis and *A. otus* also feed mostly on murine rodents (Tab. 3; Kulaeva 1977; Lundberg 1979; Mikkola 1983; Shibnev 1989; Nechaev 1991; 2003; Shepel 1992; Pukinskii 2003; 2005; Shokhrin 2005a, b; 2009). However, these owls sometimes also hunt bats (Kulaeva 1977). Remains of *M. hilgendorfi* were found in the pellets of *S. uralensis* from Sakhalin Island (4.8% of all owl prey; Nechaev 1991). Bats were recorded in the diet of the Ural owl in the valley of the River Bikin, Primorskii Region (2.0% of owl preys; Shibnev 1989) and in some areas of Kyrgyzstan (Eremchenko and Toporova 1975). Bats occurred in the diet of *A. otus* from the Perm Region (0.19%; Shepel 1992) and Sakhalin Island (0.5%; Nechaev 1991). Remains

of *Pipistrellus abramus* and *Myotis macrodactylus* were found in the pellets of the long-eared owl in Japan (Kawaguchi and Yamamota 2003; Chiba et al. 2005). We found bat remains only in spring pellets of both *A. otus* and *S. uralensis* (Tab. 2). Those of *A. otus* contained remains of *Murina hilgendorfi* and *M. ussuriensis*. To prey on ground insects (Panyutin 1974), *Murina* species mostly fly close to the earth's surface (Rossina 2007), where they may represent an easy prey for owls. Most pellets were collected in anthropogenic habitats next to Lazo village (Tab. 2) but also on the shores of Petrov Island and in the valley of the River Kievka.

The pellets of *S. uralensis* included solely bones of *M. ussuriensis*. All pellets were collected beside one nest near Mis Hill, in the valley of River Kievka. Although *M. ussuriensis* is considered rare in the Far East (Abramov et al. 2002), it was abundant in

spring pellets (Tab. 2).

CONCLUSIONS

Bats do not play a major role in the diet of owls in Southern Sikhote Alin. However in autumn and spring some bat species may form part of the owls' diet. The small number of both predator- and prey species involved probably reflects the features of owl hunting strategies and ecological and behavioural characteristics of preyed bats. Usually large and medium-sized bats living in big colonies fall victim to flying predators most frequently (Rossina et al. 2006; Shokhrin and Rossina 2008). Among them, young animals are most easily preyed upon during bat seasonal migrations.

ACKNOWLEDGEMENTS

We are indebted to V. Lebedev (Zoological Museum of Moscow State University, Moscow, Russia) for his valuable suggestions about the statistical analyses. The study was supported by the Presidium of the Russian Academy of Sciences, Program "Biosphere Origin and Evolution of Geo-Biosystems"; the Russian Foundation for Basic Research (project no. 08-04-00483-a) and by the John D. and Catherine T. MacArthur Foundation (no. 01-68420-000).

REFERENCES

- Abramov V.K., Kostenko V.A., Nesterenko V.A., Nikolaev I.G., Pikunov D.G., Sobolevskii E.I., Tiunov M.P., Yudin V.G. 2002. Mammals - Mammalia [Mlekopitayushie – Mammalia]. In: The List of the objects of the Red book of the Primorskii Region [Perechen' ob'ektov rastitel'nogo i zhivotnogo mira, zanesennih v Krasnyu knigu Primorskogo kraja]. Apostrof, Vladivostok 2002, pp. 38-39. [in Russian, without English summary].
- Aho J. 1964. The autumn food of *Asio flammeus* Pontopp. in the vicinity of the city of Tampere, South Finland. *Annales Zoologici Fennici*. Vol. 1(4): 375-376.
- Belopol'skii L.O. 1955. Birds of the Sudzukhinskii Reserve [Ptici Suhundinskogo zapovednika]: Part 2. Proceeding of the Zoological Institute, Academy of Sciences of the USSR. Moscow-Leningrad: Academy of Sciences of the USSR, Vol. 17: 225-265. [in Russian, without English summary].
- Chiba A., Onojima M., Kinoshita T. 2005. Prey of the Long-eared Owl *Asio otus* in the suburbs of Niigata City, central Japan, as revealed by pellet analysis. *The Ornithological Society*. Vol. 4: 169-172.
- Eremchenko V.K., Toporova V.I. 1975. About winter feeding of the long-eared owl in agricultural landscapes [O zimnem pitanii ushastoi sovy v usloviyakh kul'turnogo landshafta]. *Izvestiya Akademii Nauk Kirghiz. SSR*, no.5. pp. 67-68. [in Russian, without English summary].
- Kawaguchi S., Yamamota T. 2003. Food item found in pellets of Long-eared owls wintering in Ehime, Japan. *Japanese Journal of Ornithology* Vol. 52 no. 1. pp. 29-31. [in Japanese, with English summary].
- Khokhryakov S.A., Shokhrin V.P. 2002. Amphibians, reptiles, birds and mammals of the Lazovsky State Nature Reserve (Primorskii Region, Russia): Annotated lists of species. Münster: Institute of Landscape Ecology at the University of Münster. 60 p. [In Russian, in English].
- Kolomiitsev N.P., Poddubnaiya N.Ya. 1985. Data on the biology of the eagle owl *Bubo bubo* (L.) in the Lazovsky State Nature Reserve [Materialy k biologii filina - *Bubo bubo* (L.) v La-

- zovskom Zapovednike]. In: Rare birds of the Far East [Redkie ptici Dalnego Vostoka]. Vladivostok: Far Eastern Branch of the Academy of Sciences of the USSR. pp. 81-84. [in Russian, without English summary].
- Kowalski K. 1995. Taphonomy of bats (Chiroptera). *Geobios*, M.S. 18. pp. 251-256
- Kulaeva T.M. 1977. Order Strigiformes [Otryad sovoobraznie Strigiformes]. In: Birds of the Volzhsko-Kamskii Region [Ptici Volzsko-Kamskogo kraja]. Moscow: Nauka. pp. 239-257. [in Russian, without English summary].
- Kuzyakin A.P. 1950. Bats [Letuchie mishi]. Moscow: Sovetskaya Nauka. 443 p. [in Russian, without English summary].
- Lundberg A. 1979. Ecology of owls (Strigidae), especially the Ural Owl *Strix uralensis* Pall. in Central Sweden. Abstracts of Uppsala Dissertations from the Faculty of Science, no. 507. pp. 1-16.
- Mikkola H. 1983. Owls of Europe. Buteo Books. 475 p.
- Nechaev V.A. 1991. Birds of Sakhalin Island [Ptici ostrova Sakhalin]. Vladivostok: Far Eastern Branch of the Academy of Sciences of the USSR. 748 pp. [in Russian, without English summary].
- Nechaev V.A. 2003. New data on nesting of the long-eared owl *Asio otus* in the Primorskii Region [Novye dannie o gnezdovanii ushastoi sovy *Asio otus* v Primorskom kraje]. Russian ornithological journal. Express-release no. 234. pp. 958-962. [in Russian, without English summary].
- Nowak R. M. 1994. Walker's bats of the world. Johns Hopkins University Press. 287 p.
- Obuch J. 1989. Chiropteran thanatocenoses in rocky fissures. In: Hanak V., Horacek I., Gailser J. (eds.): European Bat Research 1987. Charles University Press, Praha. pp. 453
- Panyutin K.K. 1974. On the biology of *Murina leucogaster* Milne-Edwards [K biologii *Murina leucogaster* Milne-Edwards]. Proceeding of the 1st all-union bat conference [Materialy pervoi vsesoyuznoi konferentsii po rukokrylym]. Leningrad. pp. 111-112. [in Russian, without English summary].
- Priklonskii S.G. 1993. Marsh owl [Bolotnaya sova]. In: Birds of Russia and adjacent regions [Ptitsy Rossii i sopredel'nykh regionov]. Moscow. pp. 313-325. [in Russian, without English summary].
- Pukinskii U.B. 1977. Owl life [Zhizn' sov]. In: Life of our birds and mammals [Zhizn' nashikh ptic i zverey]. Vol. I. Leningrad: LSU, 240 p. [in Russian, without English summary].
- Pukinskii U.B. 2003. Nesting life of birds of the Bikin river basin [Gnezdovaya zhizn' ptits basseina reki Bikin]. Proceeding of Naturalist Society of St. Petersburg. Series 4. Vol. 86. St. Petersburg, 316 p. [in Russian, without English summary].
- Pukinskii U.B. 2005. Ural owl *Strix uralensis* (Pallas, 1771) [Dlinnokhlostaya neyasyt' *Strix uralensis* (Pallas, 1771)]. In: Birds of Russia and adjacent regions: Strigiformes, Caprimulgiformes, Apodiformes, Coraciiformes, Upupiformes, Picariae [Ptitsy Rossii i sopredel'nykh regionov: Sovoobraznie, Kozodoobraznie, Raksheobraznie, Udodoobraznie, Dyatloobraznie]. Moscow: KMK Scientific Press Ltd. pp. 72-85. [in Russian, without English summary].
- Rossina V.V. 2005. Problems of protection and monitoring of bats (Chiroptera, Mammalia) in the Lazovsky State Nature Reserve (Southern Primorskii Region) [Problemy okhrany i izucheniya rukokrylykh (Chiroptera, Mammalia) v Lazovskom zapovednike (Yuzhnoe Primor'e)]. In: Scientific research of

- natural assemblage of the Lazovsky State Nature Reserve (Proceeding of the Lazovsky State Nature Reserve), Vol. 3. Vladivostok: Russkii Ostrov, 2005. pp. 253-258. [in Russian, with English summary].
- Rossina V.V., Baryshnikov G.F., Woloszyn B.W.. 2006. Dynamics of the Pleistocene bat fauna from the Matuzka Paleolithic site (Northern Caucasus, Russia) (Chiroptera). *Lynx (Praha)*, n. s. 37 (2006). pp. 229–240.
- Rossina V.V. 2007. Bats (Chiroptera, Mammalia) of the Lazovsky State Nature Reserve and adjacent areas (Southern Primorskii Region) [Rukokrylye (Chiroptera, Mammalia) Lazovskogo zapovednika i blizlezhashchikh territorii]. *Plecotus et al.* no. 10 (2007). pp. 66-80. [in Russian, with English summary].
- Ruprecht A.L. 1990. Bats (Chiroptera) in the food of owls in the Nadnotecka Forest. *Przegląd zoologiczny*, no. 34. pp. 349-358. [in Polish, with English summary].
- Ruprecht A.L. 2005. Some aspects of myself research on bats (Chiroptera) of Poland in 1964-1990 // *Leśne Prace Badawcze*, 2005, no. 2. pp. 107-119.
- Shepel A.I. 1992. Birds of prey and owls of the Kama Region near Perm [Khishchnye ptitsy i sovyi Permskogo Prikam'ya]. Irkutsk: Irkutsk State University, 296 p. [in Russian, without English summary].
- Shibnev U.B. 1989. On the biology of the Ural owl in the Primorskii Region [O biologii dlinnokhvostoi neyasyti v Primor'e]. The newsletter of Society of naturalists of Moscow. Department of Biology. Vol. 94. Issue 5. pp. 15-25. [in Russian, without English summary].
- Shokhrin V.P. 2005a. Data on feeding of four species of Strigiformes of South-eastern Sikhote Alin [Materialy po pitaniyu chetyrekh vidov sovoobraznykh (Strigiformes) yugo-vostochno Sikhote-Alinya]. Scientific research of natural assemblage of the Lazovsky State Nature Reserve (Proceeding of the Lazovsky State Nature Reserve). Vol. 3. Vladivostok: Russkii Ostrov. pp. 240-252. [in Russian, without English summary].
- Shokhrin V.P. 2005b. Ural owl in the South-east of the Primorskii Region [Dlinnokhvostaya neyasyt' na yugovostoke Primor'ya]. Abstracts of VII Far Eastern Conference of Reserve management and studies. Birobidzhan, October 18-21, 2005. Birobidzhan: Institute for Complex Analysis of Regional Problems, Far Eastern Branch, Russian Academy of Science. pp. 298-301. [in Russian, without English summary].
- Shokhrin V.P. 2009. Biology of owls of the South Sikhote Alin [Biologiya sov Yuzhnogo Sikhote-Alinya]. In: *Owls of North Eurasia: ecology, spatial and biotopical distribution [Sovy Severnoi Evrazii]*. Moscow, 2009. pp. 246-265. [in Russian, with English summary].
- Shokhrin V.P., Rossina V.V. 2008. Food chains of owls (Strigiformes, Aves) and bats (Chiroptera, Mammalia) of South Sikhote Alin [Troficheskie svyazi sovoobraznykh (Strigiformes, Aves) i rukokrylykh (Chiroptera, Mammalia) Yuzhnogo Sikhote-Alinya]. In: *Fauna of mountain regions [Zhyvotnyi mir gornikh territorii]*. Moscow, KMK Scientific Press Ltd. pp. 513-518. [in Russian, with English summary].
- Stubbe M., Ansoerge H., Piechocki R., Lance U., Samjaa R., Bartuu D. 1989. Weitere Beiträge zur Ernährung des Uhus *Bubo bubo* (L., 1758) in der Westmongolei. *Erforschung der Biologischen Ressourcen der MVR, Halle (Saale)*. 1989 (6). pp. 25-32.
- Vorob'ev K.A. 1954. Birds of the Ussuriiskii Region [Ptitsy Ussuriiskogo kraja]. Moscow: Academy of Sciences of the USSR, 359 p. [in Russian, without English summary].

Predation on bats by owls in Russia

- Voronov G.A., Zdorikov A.I. 1988. Fish owl – *Ketupa blakistoni* Seebohm on Island Kunashir [Ribnyi filin – *Ketupa blakistoni* Seebohm na ostrove Kunashir]. In: Rare birds of the Far East and their protection [Redkie ptitsy Dal'nego Vostoka i ikh okhrana]. Vladivostok: Far Eastern Branch of the Academy of Sciences of the USSR. pp. 23-28. [in Russian, without English summary].
- Zhitkov B.M., Buturlin S.A. 1906. Data on the avifauna of the Simbirsk Region [Materialy dlya ornitofauny Simbirskoi gubernii]. The transactions of Russian Geographical Society. Vol.41. no.2. St. Petersburg. 275 p. [in Russian, without English summary].